

Evaluation of Heavy Metal Pollution in Edible Larvae of African Palm Weevil (*R. phoenicis*) in Bayelsa State, Nigeria

Thomas, C.N*; Inyang, I.R and Uwoubengha, E.R

Department of Biological Sciences, Niger Delta University,
Bayelsa State, Nigeria

*corresponding author: Email: thomascommander9@gmail.com

Abstract

Bioaccumulation of some heavy metals and selected essential elements: Mercury (Hg), Lead (Pb), Cadmium (Cd), Copper (Cu), Zinc (Zn) and Manganese (Mn) was investigated in the edible larvae of African Palm Weevil (APW) (Rhynchophorus phoenicis) from polluted sites and unpolluted site. Cadmium pollution was confirmed by the presence of 0.025 mg/100g to 0.037mg/100g of cadmium in the larvae, above the limit of 0.02mg/100g was hazardous to health of consumers. Higher concentration of manganese (1.804 mg/100g- 2.848mg/100g) found in the larvae was a risk to the health of consumers because it was above the limit of 0.05mg/100g. There was low concentration of zinc (2.79mg/100g-2.89 mg/100g) and copper (0.07.40mg/100g-2.89 mg/100g) below the limit of 9.94mg/100g. There was no bioaccumulation of lead and mercury in the larvae because it contained trace amount of 0.0001mg/100g of both heavy metals, which was below the limit of 0.0006 mg/100g. Larvae of R. phoenic are good biomarkers for monitoring heavy metal pollution in the environment. Effective control and management of toxicity of cadmium and manganese is required to save humans and biota in the affected communities and the environment.

Key words: Bioaccumulation, heavy metals, and *R. phoenicis*, larvae

1. Introduction

The larvae of the African Palm weevil (*Rhynchophorus phoenicis*) are popularly known as “diet” in Warri and “Suya” in Bayelsa State. It is a highly cherished traditional delicacy in many parts of Nigeria and other countries in Africa (Thomas, 2003 and Ekpo, 2003). (Defoliart, 1992; Allotey and Mpuchane, 2003, Choon-Fah *et al*, 2008). The Larvae of *R. phoenicis* are often hawked along major roads and markets across the Niger Delta States of Nigeria (Ekrakene and Igeleke, 2007). The mode of preparation of the larvae for eating differs from one ethnic group to another (Thomas *et al*, 2017). It is occasionally consumed in raw form roasted or fried but the fried form was more acceptable in the Niger Delta where it is usually eaten with tapioca or bread by the Urhobo in Delta State, whereas the Ijaw people eat it with soaked garri, farina and any staple food (Thomas 2003; Thomas, *et al*, 2005). The larvae are endowed with lot of nutrients, hence pregnant women eat it as a source of essential nutrients (Ekpo, 2003). The use of the larvae was beyond the nutritional value, as it was believed to possess some medicinal properties, hence the Itsekiri people in Delta State use the life larvae to cure some ailments in infants. It was also recommended for traditional healing of new born babies in some Ikwerre communities in Rivers State. A survey of its food uses showed that the larvae of *R. phoenicis* have different names in different ethnic groups, tribes and states in Nigeria, which confirms its widespread utilization as an important food in different parts of the oil rich palm belt, which include Bayelsa, Rivers, Cross River, Akwa

Ibom, Delta, Edo, Imo, Benue, Anambra and other states in southern Nigeria (Thomas 2003; Thomas and Briyai, 2019). Incidentally, the palm belt span across the oil-rich geological zone which equally produces commercial crude oil. Therefore, the larvae of the palm weevil had been associating with heavy metals released to the environment; hence, conscious effort is required to protect the masses from the intake of harmful pollutants, especially long-term exposure to the heavy metals. This study was carried out to ascertain the amount of some heavy metals in the edible larvae of African palm weevil in some impacted locations in Bayelsa State, where crude oil exploration had continued since 1956 till date

2. Materials and Method

2.1 Description of study area

The Gbarain gas gathering project location at Obunagha community in Gbarain Kingdom, Yenagoa local Government area was set as site 1 sitting on a coordinate of Longitude 4.9514°N, and Latitude 6.3492°E. Site 2 was set at Etelebou flow station at Ogboloma Community in Gbarain Kingdom sitting on a Longitude of 5.0440 °N and Latitude of 6.3379 °E, while Site 3 was taken at Tombia round about along Imiringi road within Yenagoa municipality sitting on a Longitude of 4.9514°N and Latitude 6.3492°E. The control site (a non-polluted area) was taken at Toru-Ebeni forest in Sagbama Local Government Area sitting on a coordinate of Longitude 4.731°N and Latitude of 6.1090°E all in Bayelsa State of Nigeria (Fig. 1).

2.2 Source of Materials

The larvae of *R. phoenicis* were obtained from infested raphia palms from the four locations (Fig. 1). In each site, three matured raphia palms (*Raphia hookeri*) that has been tapped for wine was cut down and left for over eight to twelve weeks on the ground at the various sites, after which it was harvested by cutting - open the infested portions of the trunk with an axe and matchete. The larvae were extracted manually from the tissues of the decaying palms and transferred in aerated plastic buckets with some quantity of the fibrous tissues and transported to the laboratory for analysis.

2.3 Heavy Metals Analyses Procedures

Samples were oven dried at 80-90°C and was grinded with a mortar pestle, 2g of the sample was weighed into a glass beaker (100ml); 5g of sample was transferred into 100ml glass beaker, while a mixture of 2ml concentrated nitric acid (HNO₃), 10ml of concentrated hydrochloric acid (HCl) and about 20ml distilled water was added. Samples were digested on a coming PC-351 model hot plate at medium to low heat until about 5ml concentrated extract was left. The content of the beaker was left to cool for about 30 minutes. The sample solution was filtered and quantitatively transferred into 50ml standard volumetric flask. Finally, the filtered solutions were marked up to the 50ml graduation line using distilled water. The filtrate was used in the determination of the following elements (Zn, Pb, Cd, Ca, Mn and Hg) using the GBC 908PBMT model flame atomic absorption spectrophotometer (FAAS). The total metal concentration of each sample was reported in units of ppm (mg/100g) and converted to mg/100g. The results obtained are the means of triplicate samples with standard deviations.

2.4 Statistical Analysis

The version 23 of SPSS was adopted for the statistical analysis. Analysis of variance (ANOVA) was used for mean separation, while Duncan multiple range was adopted to establish the degree of significant differences in the results from the various sites.




Fig 1: A map (B) showing the study area and site locations depicted in 4 distinct colours as extracted from the map of Nigeria (A).

- Site 1: Gbarain gas gathering project site, Obunagha in Gbarain Kingdom in Yenagoa Local Government Area
- Site 2: Etelebou flow station, Ogboloma in Gbarain Kingdom in Yenagoa Local Government Area
- Site 3: Gas station, Tombia round about along Imiringi road in Yenagoa municipality
- Site 4: Control Site (Non-Polluted site at Toru Ebeni forest in Sagbama Local Government Area).

3.0 Results

The results (Table 1) showed that there was no bioaccumulation of mercury in the larvae from all the sites because the amount of mercury (0.0001 mg/100g) was below the standard limit (0.0006 mg/100g) with no significant difference ($p>0.05$) in all the sites. The amount of Lead (0.0001 mg/100g) was below the standard limit (0.0001 – 0.0003 mg/100g) (Harvard, 2018) in all the sites, with no significant difference ($p>0.05$) when compared to the control site.

The amount of copper in the larvae ranged between 0.074 – 2.879 mg/100g was below the standard limit of 9.94 mg/100g. There was significant difference in the amount of copper in the polluted sites, which was higher than the control site ($p<0.05$). Similarly, the amount of zinc in the larvae (2.790 – 2.890 mg/100g) was below the WHO acceptable limit of 9.940 mg/100g (Harvard, 2018). Incidentally, the results showed that the larvae of *R. phoenic* is contained low amounts of cadmium (0.025 – 0.037 mg/100g) from the control and polluted sites respectively, was above the permissible limit of 0.02 mg/100g (Harvard 2018), with no significant difference ($p>0.05$). The amount of manganese in the larvae ranged between 1.804 – 2.862 mg/100g from the control site and polluted was above the WHO permissible amount of 0.05 mg/100g (Harvard, 2018). There was no significant difference in the amount of manganese in the larvae from the polluted sites ($p>0.05$), but the amount of manganese in the larvae from the control site was significantly lower than the amount of manganese in the larvae from the polluted sites.

Discussion: The findings from this study revealed the presence of higher concentration of manganese which ranged between 1.804mg/100g to 2.862mg/100g in *R. phoenicis* larvae was beneficial to consumers, because manganese is an essential element required for bone formation and strengthening of weakened bones in people who suffers from osteoporosis (WHO, 1977). In addition, manganese is a co-factor in the active sites of many enzymes involved in normal development, maintenance of nerves and immune cell functions, regulation of blood sugar and vitamins among others (Antonini *et al* 2006, Bailey *et al*, 2017 and Ben-shahar, 2018).

However, when people are chronically exposed to manganese, it become a risk factor due to its toxic effect on many organ systems (“typical parkinsonian syndrome” or “manganism”) arising from the accumulation of manganese in the brain. The above pathological condition was attributed to the presence of dopamine, GABA and glutamate neural signaling in insect brain which affects the foraging activities of pollinators (honey bees) (Ben-shahar, 2018). The presence of 2.79mg/100g – 2.80mg/100g of zinc in the larvae of *R. phoenicis* was of great nutritional importance because zinc is an essential element that is rarely present in diets of vegetarians. Therefore, vegans can be advised to consume the larvae of *R. phoenicis* as supplementary source of zinc to ameliorate chronic zinc deficiency. Zinc is required in the active-sites of about 300 enzymes involved in metabolizing glycosides and proteins, as well as insulin secretion by the pancreas, to enhance immunity, resolve sexual infertility, improve growth and cause faster rate of healing of wounds.

Table 1: Heavy Metals Composition in *R. phoenicis* larvae from polluted and Control sites.

Parameters (mg/100g)	Polluted site 1 (OGGP)	Polluted site 2 (EFS)	Polluted site 3 (IMR)	Control site 4 (TE)	WHO limits (Harvard, 2018)
Mercury	0.0001±0.0 ^a	0.0001±0.0 ^a	0.0001±0.0 ^a	0.0001±0.0 ^a	0.0006
Lead	0.0001±0.02 ^c	0.0001±0.02 ^c	0.0001±0.02 ^c	0.0001±0.02 ^c	0.0001-0.003
Copper	0.879±1.7 ^b	0.910±1.7 ^c	2.89±3.1 ^d	0.07.40±1.6 ^a	9.94
Cadmium	0.037±0.4 ^a	0.025±0.3 ^a	0.037±0.4 ^a	0.025±0.9 ^a	0.02
Zinc	2.80±3.1 ^a	2.79±3.0 ^a	2.89±3.1 ^a	2.790±3.0 ^a	9.94
Manganese	2.848±3.1 ^b	2.831±3.1 ^b	2.862±3.1 ^b	1.804±2.5 ^a	0.05

Different superscript (a,b,c,d) along row indicates a significant different among site compared to the control. ($p < 0.05$)

Key:

Site 1: OGGP – Obunagha –Gas gathering project site

Site 2: EFS – Etelebou Flow Station

Site 3: IMR – Imiringi Road, Yenagoa.

Site 4: Control – Toru – Ebeni bush

Zinc is required in the active-sites of about 300 enzymes involved in metabolizing glycosides and proteins, as well as insulin secretion by the pancreas, to enhance immunity, resolve sexual infertility, improve growth and cause faster rate of healing of wounds. Prolonged zinc deficiency was often associated with liver disease and pernicious anemia (ICMSF, 1986).

Cadmium usually accumulates in the roots of terrestrial plants and bind to the cell wall, through which it is easily ingested by insects (grasshoppers) and aquatic invertebrates (especially molluscs) which feed on the plants for growth and reproduction. Similarly, in the case of the adult palm weevils, it lay its eggs inside the feeding holes which later hatch and develop through its life cycle in the tissues of an infested raffia or young oil palm (Kehinde *et al*, 2017). Atmospheric cadmium emitted to the air through gas flaring, deposits sparingly on the soil and absorbed through the roots of plants which are eaten by insects and other animals. Worst, in recent times, is the menace of illegal refining of crude oil obtained from vandalized pipelines by unemployed youths, as their means of survival, thereby contributing to the accumulation of cadmium and other heavy metals in the air and soil of Niger Delta (NCMCR, 2003). The frequent occurrence of crude oil spills, uncontrolled incineration of wastes at dumpsites, bush-burning by peasant farmers and occasional use of chemical fertilizers are additional sources of heavy metal emissions into the air (AMAP, 1998). Although, cadmium is not useful to plants and animal life, low concentrations are tolerated by binding it to special proteins to make it harmless. When it exceeds the permissible level of 0.02mg/100g, it affects the health of people by causing kidney damage and lung emphysema in humans (IPCS WHO, 1992a and 1992b).

The International Agency for Research in cancer confirmed that cadmium is one of the most carcinogenic elements in humans as it frequently causes lung cancer and occasionally causes prostate cancer in people who are exposed to it by their occupation (Waalkes, 2000 and IARC, 1993). The results of this study have revealed that consistent bioaccumulation of heavy metals is taking place in the soils of the impacted areas due to systemic accumulation of pollutants emitted through gas flares from crude oil flow stations at Etelebou, plus pollutants from the Gbarain-Gas gathering plant negatively affects the immediate host communities including: Obunagha, Okolobiri, Ogboloma, Koroama, Nedugo-Agbia, Polaku and all the neighbouring communities of Ekpetiama clan, especially Gbarantoru, Tombia and Akaibiri in Yenagoa local government area of Bayelsa State. The negative impact spreads over Imiringi road axis where massive oil exploration activities had continued at flow stations which flares gases in Ogbia local government area where the first commercial oil well was discovered at Oloibiri in 1957. The results further revealed that Toru-Ebeni which was taken as a control site due to the non-existence of oil exploration activities, yet it suffered heavy metal pollution through anthropogenic sources (humans, animals, insects) and by natural geochemical actions of wind, water and other abiotic factors (Lee *et al*, 2006). This implied that, the entire Bayelsa State is affected by chemical pollution by extension of the dispersion of the pollutants through air, surface and ground water resources (Ukponget *al*, 2013). It had been reported that heavy metal accumulation causes growth inhibition and reduces the fecundity and hatchability of eggs in insects. This means that bio-accumulation of cadmium and other heavy metals would affect the reproductivity of *R. phoenicis* which is a nutritious edible insect species that enhances the socio-economic wellbeing and livelihood of rural communities in the Niger Delta of Southern Nigeria, and other African developing economies. Therefore, it calls for intervention by international agencies that are concerned with food security, social health and environmental protection to respond with finding solution to the environmental menace of heavy metal pollution in Bayelsa state and indeed, the Niger Delta of Nigeria.

Conclusion: The findings of this study revealed that, the larvae of *R. phoenicis* contained significant amounts of cadmium (0.025-0.037mg/100g) which was above the permissible limit of 0.02mg/100g. Therefore, cadmium pollution was confirmed in the edible larvae of *R. phoenicis*, which exposes the rural communities to grave health hazards, arising from the carcinogenicity of cadmium which causes lung cancer and occasionally responsible for prostate cancer. So, all concerned authorities should respond to control the environmental threats to life of the people in the affected communities. The edible larvae of *R. phoenicis* also contained high concentration of manganese which ranged between 1.84mg/100g to 2.83mg/100g which exceeded the permissible limit of 0.05mg/100g of manganese. Therefore, further studies on the impact of environmental manganese on the biology and physiology of *R. phoenicis* as well as, its role on the ecological balance of the food web of the ecosystem should be conducted without delay. The edible larvae of *R. phoenicis* should be used as biomarkers of heavy metal contamination during environmental monitoring and impact assessment. There is urgent need to adopt effective control measures for cadmium and manganese pollution to avert the grave consequences of the affected host communities of Gbarain, Ekpetima and Epie-Atissa clans in Yenagoa Local Government Area, and indeed the whole Bayelsa State from imminent danger of chemical pollution. The concerned authorities of government should support studies on how to prevent further increase to hazardous levels in the accumulation of copper and other heavy metals, to save the environment and its inhabitants from disaster in Bayelsa State and the entire Niger Delta Area of Nigeria, where oil exploration and illegal refining activities are rampant in recent times.

References:

- Allotey J and Mpuchane S (2003): Utilization of useful insects as food source. *African Journal of Food, Agriculture, Nutrition and Development*, 3:1-8.
- AMAP (1998): Assessment Report; Arctic pollution issues. Arctic Monitoring and Assessment Programme, Oslo.
- Antonini, J.M; Santamaria, A. B; Jenkins, N.T; Albini E and Lucchini R (2006): Fate of manganese associated with the inhalation of welding fumes: potential neurological effects. *Neurotoxicology* 27,304 – 310 DOI: 10.1016/j.neuro.2005.09.00
- Bailey, L.A; Kerper, L.E and Godman, J.E (2017): Derivation of an occupational exposure level for manganese in welding fumes. *Neurotoxicology* 64, 166 – 167. DOI: 10.1016/J.neuro.2017.06.009.
- Ben-shahar Yehuda (2018): The impact of Environmental Manganese Exposure on insect Biology. *Frontiers in Genetics: Mini-Review*. Published online www.frontiersin.org.doi: 10.3389/fygne.2018.00070 (Pp. 1-7)
- Choon-Fah J.B, Chin-Chin E, Pang-Hung Y. and Amartalington R. (2008): Growth Performance of the red-striped weevil, *Rhynchophorus schach* Oliv.(Insecta; Coleoptera; Curculionidae) on meridic diets. *Am. J. Agric. Biol.Sci*; 3:403-409
- Defoliart G. R. (1992): A concise summary of the general nutritional value of insects. (<http://www.food.insects.com>) Insects as Human Food
- Ekpo K.E (2003): Biochemical investigation of the Nutritional value and toxicological safety of Entomophagy in southern Nigeria. Ph.D. thesis
- Ekrakene T. and Igeleke C.L (2007): Microbial Isolates from the roasted larvae of the palm weevil (*Rhynchophorus phoenicis* (F) from Edo and Delta state of Nigeria. *Am.J.Biol. Applied Sci*, 1:763 -768.
- Harvard Medical School (2018), Harvard Health Letter, Published July, 2018: Recommended Daily intake of Vitamins and Minerals. <https://www.lenntech.com/recommended-dailyintake.html#ix226TgkNITJS>

- I.C.M.S.F (1978): Micro Organisms in Food: sampling for the microbial analysis. Principles and Specification Blackwell Publications: New York, U.S.A
- IARC (1993): Cadmium and Cadmium Compounds (Group 1), IARC monographs Vol.58: 1993. *International Agency for Research on Cancer*, Lyon.
- Kehinde Ayeni, OluwafemiAjayi and Dolapo Odeyemi (2017). Studies on Heavy Metal Accumulation in plants, insect and soil in public dumpsite in Ado-Ekiti, Ekiti state. *International Journal of Sciences: Basic and Applied Research (IJSBAR)*. 36(2):18-26
- Lee C, Li S-L, Shi X.W and Cheung S.C (2006): “Metal Contamination in Urban, Suburban and Country Park soils of Hong Kong: a study based on GIS and multivariate statistic” *Science of the Total Environment*, Vol. 356, PP. 45 -61
- NCMCR (2003): Nordic Council of Ministers, Cadmium Review. Report: 1(4):1-24, January, 2003.
- Thomas C.N and Briyai F.O (2019): Traditional Consumption of *Rhynchophorusphoenicis* (F.) as human food in Niger Delta Area, Nigeria. *Journal of Biology and Genetic Research* (JBGR) ISSN 2545 – 5710.
- Thomas, C.N; Okwakpam, B.A; Empere, C.E and Ogbalu, O.K (2005) Utilization of *Rhynchophorus phoenicis* F. (Coleoptera: Curculionidae) as Human food in Niger Delta Area of Nigeria. *Niger Delta Biologia*. Vol. 6 (2): 18-22.
- Thomas Commander N. (2003): Biology, Utilization and Rearing of the African Palm weevil (*Rhynchophorusphoenicis* F.) on palms of the Niger Delta. Rivers State University, Nkpolu, Port-Harcourt, Nigeria Ph.D. thesis.
- Ukpong E.C, Antigha R.E and Moses E.O (2013): Assessment of Heavy Metals content in Soils and Plants around Waste Dumpsite in Uyo Metropolis, Akwa-Ibom state. *International Journal of Science and industrial Research*, vol.51 (1), PP. 329 -332.
- Waalkes, M.P (2000): Cadmium carcinogenesis in review. *Journallnorganic Bio-chemistry* 7a: 240 – 244.
- WHO (1992b): Cadmium – Environmental aspects. Environmental Health criteria 135. World Health Organization, International Programme on Chemical Safety (IPCS), Geneva, Switzerland.
- WHO 1992a. Cadmium. Environmental Health Criteria 134. World Health Organization, International Programme on Safety (IPCS), Genera, Switzerland.