

Phytochemical Variability in Two Cocoyam Varieties Grown in Port Harcourt

Victor I. Ajie¹, Wumi Esther Adelegan²

¹ Department of Plant Science and Biotechnology, Rivers State University, Npkolu-Orowuroko, Port-Harcourt, Rivers State, Nigeria.

² Department of Science Laboratory Technology Federal Polytechnic Ugep, Cross rivers State
DOI: 10.56201/rjps.t.vol.8.no1.2025.pg39.46

Abstract

The comparative analysis of the phytochemical profiles of Colocasia esculenta (L.) Schott and Xanthosoma sagittifolium, two underutilized tuber crops with potential nutritional and medicinal applications, was studied. The research was carried out at the Department of Plant Science and Biotechnology, University of Port Harcourt and investigation focused on determining levels of saponins, tannins, alkaloids, flavonoids and oxalates in both species. Results revealed that Colocasia esculenta had higher concentrations of saponins (0.682%), tannins (0.0001235%), and alkaloids (1.065%), indicating it's potential for stronger therapeutic benefits, especially in traditional medicine. In contrast, Xanthosoma sagittifolium exhibited higher flavonoid content (2.76%), suggesting superior antioxidant properties, though it also had a higher oxalate content (0.558%). The study concludes that both plants offer valuable contributions to health, with Colocasia esculenta being more suitable for medicinal purposes and Xanthosoma sagittifolium for antioxidant intake. It is recommended that the use of these plants based on their phytochemical strengths is promoted and additional research on the bioavailability and safety of their compounds be conducted.

Keywords: Cocoyam varieties, Phytochemical composition, Comparative analysis, Port Harcourt, Bioactive compounds

1. Introduction

Cocoyam is a monocotyledonous, herbaceous, perennial and vegetatively propagated plant of the family Araceae. This family is characterized by its morphological diversity, the presence of many forms of calcium oxalate crystals, and flowers with a spadix of small, bisexual, or unisexual flowers, covered by a spathe (Henriquez *et al.* 2014; Matthews 1995). In Nigeria, the term "cocoyam" pertains to two varieties of edible aroid plants, specifically *Colocasia esculenta* (L.) Schott and *Xanthosoma sagittifolium*. Africa is the major producer of cocoyam, with West and Central Africa, notably, Nigeria, Ghana, and Cameroon contributing to over 60% of the total African production (Onyeka, 2014). According to Falade and Okafor (2014), Nigeria is the largest producer of cocoyam in the world with an annual production of 3.450 million metric tonnes in 2012, representing 72.2 %, 57.7 % and 45.9 % of total production in West Africa, Africa and the World respectively. Cocoyam is a rich source of carbohydrates and serve as a staple food for over 400 million people in the tropics (Djeugap *et al.*, 2014).

Colocasia esculenta and *Xanthosoma sagittifolium* are two distinct, yet closely related root crops belonging to the Araceae family, both of which are widely cultivated in tropical and

subtropical regions (Gelleh *et al.*, 2018). *Colocasia esculenta* is typically characterized by its large, heart-shaped leaves and corms that have a smooth, brown skin with a starchy, white or purple-tinged flesh (Moorthy, 2004). In contrast, *Xanthosoma sagittifolium* has arrow-shaped leaves with corms that are generally larger, rougher, and more fibrous, with a flesh that ranges from white to yellow (Manner, 2011). Both plants are significant in traditional diets, offering a rich source of carbohydrates, dietary fiber, and essential micronutrients, making them vital for food security in many regions (Lim, 2015). Despite their similarities, they differ in their cultivation requirements, with *C. esculenta* preferring wetter, lowland conditions and *X. sagittifolium* thriving in drier, upland areas (Manner, 2011).

Phytochemicals are the plant derived chemicals possessing numerous herbal & medicinal properties. Medicinal plants with important bioactive compounds play a significant role in meeting the global healthcare needs. In fact, eighty percent of African populations use some form of traditional herbal medicine, and the worldwide annual market for herbal products approaches US\$ 60 billion (Gunjan *et al.*, 2015). Through phytochemical screening one could detect the various important compounds which could be used as the base of modern drugs that curing various diseases (Nilofer *et al.*, 2013). Cocoyam possesses a range of biologically active phytoconstituents such as flavonoids, sterols, glycosides, and other micronutrients (Prajapati *et al.*, 2011). Phytochemically, while both species contain valuable bioactive compounds such as flavonoids and phenolics, their profiles may differ, potentially leading to variations in their medicinal properties.

Colocasia esculenta and *Xanthosoma sagittifolium* are not only dietary staples in many tropical regions but also possess a wealth of bioactive compounds that could offer therapeutic benefits. The phytochemical profiles of *C. esculenta* and *X. sagittifolium* have not been extensively compared, leaving a gap in our understanding of their relative health-promoting properties. This knowledge is critical given the growing interest in plant-based compounds for their potential health benefits. Therefore, this study was carried out to compare and analyze the phytochemical profiles of corms of *Colocasia esculenta* and *Xanthosoma sagittifolium*.

2. Materials and Methods

2.1 Study Area

This study was carried out at the Department of Plant Science and Biotechnology, University of Port Harcourt, Choba, Rivers State. It is located on Latitude 4°54'42" N and Longitude 6°54'00"

2.2 Source of Sample

Colocasia Esculenta and *Xanthosoma Sagittifolium* were sourced from Aspatech Integrated Farms at Trans-Kalabari road, Rumuolumeni, Port Harcourt, Rivers State.

2.3 Sample Collection

The corms were carefully harvested, with attention to avoid contamination and transported immediately to Department of Plant science and Biotechnology, Rivers State University, located on Latitude 6°58'39.12"E and Longitude 4°48'18.50 where they were identified by the plant taxonomist Dr. (Mrs) M. G. Ajuru.

2.4 Sample Description

The *Colocasia esculenta* samples were characterized by smooth, brown skin with a starchy, white or purple-tinged flesh, while the *Xanthosoma Sagittifolium* samples exhibited generally larger, rougher, and more fibrous flesh.

2.5 Sample Preparation

Corms of *Colocasia esculenta* and *Xanthosoma Sagittifolium* were washed thoroughly to remove soil and debris. The cleaned samples were then sliced into smaller pieces and air-dried to remove moisture. Following drying, the samples were ground into fine powder using a mechanical grinder, ensuring uniform particle size for consistent analysis.

2.6 Methodology

The methods of AOAC, (2005) were used for the determination of oxalate, saponin, tannin, alkaloid and flavonoid compositions. All analysis were done in triplicate.

3. Results

3.1 Saponin Content

Colocasia esculenta had a slightly higher saponin content (0.682%) compared to *Xanthosoma sagittifolium* (0.625%).

3.2 Tannin Content

Tannin content was very low in both species, but *Colocasia esculenta* had a slightly higher tannin percentage (0.0001235%) than *Xanthosoma sagittifolium* (0.0000985%).

3.3 Alkaloid Content

There was a significant difference in alkaloid content between the two species. *Colocasia esculenta* had a much higher alkaloid percentage (1.065%) compared to *Xanthosoma sagittifolium* (0.17%).

3.4 Flavonoid Content

Xanthosoma sagittifolium had a higher flavonoid content (2.76%) compared to *Colocasia esculenta* (1.64%).

3.5 Oxalate Content

Xanthosoma sagittifolium also had a higher oxalate content (0.558%) compared to *Colocasia esculenta* (0.3685%).




Fig 1: Comparative Phytochemical Contents of *Xanthosoma Sagittifolium* and *Colocasia Esculenta*

4. Discussion

The results indicate that *Colocasia esculenta* has a slightly higher saponin content compared to *Xanthosoma sagittifolium*. Saponins are a group of naturally occurring plant glycosides known for their ability to form soap-like foams when mixed with water. They have been associated with various health benefits, including cholesterol-lowering properties, immune-boosting effects, and potential anticancer activities (El Aziz *et al.*, 2019). The presence of saponin is in agreement with the findings of McEwan *et al.* (2014). The relatively similar saponin levels in both plants suggest that they could both provide comparable health benefits related to saponin consumption.

The tannin content in both *Xanthosoma sagittifolium* and *Colocasia esculenta* was very low, with *Colocasia esculenta* having a slightly higher level compared to *Xanthosoma sagittifolium*. Tannins are polyphenolic compounds known for their astringent properties, which can contribute to the plant's defense against herbivores and pathogens (Fraga-Corral *et al.*, 2020). In terms of human health, tannins are recognized for their antioxidant properties and their potential to inhibit the growth of bacteria, viruses, and fungi (Tong *et al.*, 2022). The low tannin levels in both plants is similar to findings of Abdurashid and Agwunobi (2012) and suggest that their contributions to antioxidant activity may be limited.

A significant difference is observed in the alkaloid content, with *Colocasia esculenta* containing much higher levels compared to *Xanthosoma sagittifolium*. Alkaloids are a diverse group of nitrogen-containing compounds that are often pharmacologically active, making them important in traditional medicine (Gupta et al., 2020). They can have a variety of effects, including analgesic, anti-inflammatory, and antimicrobial activities (Jafaar et al., 2021). The presence of alkaloids is also in agreement with the findings of McEwan et al. (2014), but the higher alkaloid content in *Colocasia esculenta* suggests that it may have more potent medicinal properties, potentially offering greater therapeutic benefits. This could make *Colocasia esculenta* more valuable in contexts where alkaloid-related health effects are desired.

Xanthosoma sagittifolium exhibited a higher flavonoid content compared to *Colocasia esculenta*. Flavonoids are well-known for their antioxidant properties, which can help in preventing cellular damage caused by free radicals. They also have anti-inflammatory, antiviral, and anticancer potential (Zhang et al., 2018). Udo et al. (2020) also reported significant flavonoid content in wild cocoyam. The significantly higher flavonoid content in *Xanthosoma sagittifolium* suggests that it may offer more substantial antioxidant benefits, making it a potentially more effective dietary source for reducing oxidative stress and related diseases.

Xanthosoma sagittifolium also had a higher oxalate content compared to *Colocasia esculenta*. Eleazu et al. (2018) also reported the presence of oxalates in cocoyam corms. Oxalates are organic compounds found in many plants, and while they can have some beneficial effects, they are also known to bind with calcium to form kidney stones in susceptible individuals (Savage and Klunklin, 2018). The presence of oxalates, which is responsible for the acrid taste or irritation in its raw corm, is the major factor inhibiting the utilization of cocoyam (Oke and Bolarinwa, 2012) and the higher oxalate content in *Xanthosoma sagittifolium* suggests that caution may be needed when consuming this plant.

5. Conclusion

The comparative analysis of *Xanthosoma sagittifolium* and *Colocasia esculenta* reveals distinct phytochemical profiles that offer unique nutritional and medicinal benefits. *Colocasia esculenta* is richer in saponins, tannins, and alkaloids, indicating its potential for stronger therapeutic effects, particularly in traditional medicine. On the other hand, *Xanthosoma sagittifolium* boasts a higher flavonoid content, which may provide superior antioxidant benefits, making it valuable for reducing oxidative stress and related health issues. However, the higher oxalate content in *Xanthosoma sagittifolium* necessitates caution, especially for individuals at risk of kidney stone formation. Overall, these findings underscore the importance of understanding the phytochemical composition of underutilized plants to optimize their use in food and medicine.

It is therefore recommended that *Xanthosoma sagittifolium* should be promoted as a dietary source of antioxidants, especially for individuals seeking to enhance their intake of natural compounds that combat oxidative stress, but with caution due to its higher oxalate content. Also, *Colocasia esculenta* should be further explored for its higher alkaloid content, which could be harnessed for developing herbal remedies with potential anti-inflammatory and antimicrobial properties. Finally, additional studies should be conducted to explore the

bioavailability and safety of the phytochemicals in these plants, particularly the impact of their oxalate content on health, to ensure they are safe for widespread consumption.

References

- Abdurashid, M. and Agwunobi, L. N. 2012. Tannia (*Xanthosoma sagittifolium*) cocoyam as dietary substitution for maize in broiler chicken. *Greener Journal of Agricultural Sciences* 2(5): 167-171.
- AOAC. 2005. Official methods of analysis of AOAC international. 18th edition. Association of official analytical chemists, Washington, D.C., USA.
- Djeugap, J. F., Azia, T. A. and Fontem, D. A. 2014. Effect of Compost Quality and Microbial Population Density of Composts on the Suppressiveness of *Pythium myriotylum*, Causal Agent of Cocoyam (*Xanthosoma sagittifolium*) Root Rot Disease in Cameroon. *International Journal of Sciences: Basic and Applied Research (IJSBAR)*.
- El Aziz, M. M. A., Ashour, A. S., & Melad, A. S. G. (2019). A review on saponins from medicinal plants: chemistry, isolation, and determination. *J. Nanomed. Res*, 8(1), 282-288.
- Eleazu, C., Sampson A., Saidu, S., Eleazu, K. and Egediwe-Ekeleme, C. 2018. Starch digestibility, polyphenol contents and in-vitro alpha amylase inhibitory properties of two varieties of cocoyam (*Colocasia esculenta* and *Xanthosoma maffafa*) as affected by cooking. *Journal of Food Management and Characterisation*, 12: 1047 -1053.
- Falade, K. and Okafor, C. 2014. Physical, functional, and pasting properties of flours from corms of two Cocoyam (*Colocasia esculenta* and *Xanthosoma sagittifolium*) cultivars. *Journal of Food Science and Technology*. 52. 10.1007/s13197-014-1368-9.
- Fraga-Corral, M., García-Oliveira, P., Pereira, A. G., Lourenço-Lopes, C., Jimenez-Lopez, C., Prieto, M. A., & Simal-Gandara, J. (2020). Technological application of tannin-based extracts. *Molecules*, 25(3), 614.
- Gelleh, I. D., Okeke, U. H., Babalogbon, B. A. and Mangut, Y. S. 2018. Land suitability analysis for the production of cocoyam in Benue state, Nigeria. *Earth Sciences Malaysia*, 2(2): 25 –30.
- Gunjan, M., Naing, T., Saini, R., Ahmad, A., Naidu, J. and Kumar, I. 2015. Marketing trends and future prospects of herbal medicine in the treatment of various disease. *World Journal of Pharmaceutical Research*, vol. 4, no. 9, pp. 132–155.
- Gupta, A. K., Tulsyan, S., Thakur, N., Sharma, V., Sinha, D. N., & Mehrotra, R. (2020). Chemistry, metabolism and pharmacology of carcinogenic alkaloids present in areca nut and factors affecting their concentration. *Regulatory Toxicology and Pharmacology*, 110, 104548.
- Henriquez, C. L., Arias, T., Pires, J. C., Croat, T. B. and Schaal, B. A. 2014. Phylogenomics of the plant family Araceae. *Mol Phylogenet Evol* 75:91–102.

- Jafaar, H. J., Isbilen, O., Volkan, E., & Sariyar, G. (2021). Alkaloid profiling and antimicrobial activities of *Papaver glaucum* and *P. decaisnei*. *BMC Research Notes*, *14*, 1-7.
- Lim, T. K. 2015. Edible Medicinal and Non-Medicinal Plants. Volume 9: Modified Stems, Roots, Bulbs. New York: Springer.
- Manner, H. I. 2011. Farm and forestry production and marketing profile for Tannia (*Xanthosoma* spp.). In: *Specialty crops for Pacific Island Agroforestry*, pp. 1-16, (Elevitch, C.R., ed). Permanent Agriculture Resources, Hawaii, USA.
- Matthews, P. J. 1995. Aroids and the Austronesians. *Tropics* 4:105–126.
- McEwan, R., Shangase, F. N., Djarova, T. and Opoku, A. R. 2014. Effect of the three processing methods on some nutrients and anti-nutritional factor constituents of *Colocasia esculenta* (Amadumbe). *African Journal of Food Science*, *8*(5): 287 –292.
- Moorthy, S. N. 2004. Tropical sources of starch. p. 321-359. In Ann-Charlotte Eliasson (ed) *Starch in food: Structure, function and applications*. Woodhead Publishing Series in Food Science, Technology and Nutrition.
- Nilofer, S., Yogendra, K., Misra, A. K. and Lokho, P. 2013. Phytochemical screening to validate the ethnobotanical importance of root tubers of *Dioscorea* species of Meghalaya, North East India.; *1*(6):62-69.
- Oke, M. O. and Bolarinwa, I. F. 2012. Effect of fermentation on the chemical properties and oxalate content of cocoyam (*Colocasia esculenta*) flour. *International Scholarly Research Notices*; Article ID 978709, 4pages.
- Onyeka, J. 2014. Status of Cocoyam (*Colocasia esculenta* and *Xanthosoma* spp.) in West and Central Africa: Production, Household Importance and the Threat from Leaf Blight.
- Prajapati, R., Kalariya, M., Umbarkar, R., Parmar, S. and Navin, S. 2011. *Colocasia esculenta*: A potent indigenous plant. *International journal of Nutrition, Pharmacology, Neurological Diseases*. 1. 90. 10.4103/2231-0738.84188.
- Savage, G., & Klunklin, W. (2018). Oxalates are found in many different European and Asian foods-effects of cooking and processing.
- Tong, Z., He, W., Fan, X., & Guo, A. (2022). Biological function of plant tannin and its application in animal health. *Frontiers in veterinary science*, *8*, 803657.
- Udo, M. D., Eyoh, G. D., Jimmy, C. P., & Ekpo, U. E. (2020). Nutrient Composition, Mineral and Anti-Nutrient Components of Processed Wild Cocoyam (*Caladium bicolor*, (Ait) Vent). *Current Agriculture Research Journal*, *8*(2).
- Zhang, H., Yang, Y. F., & Zhou, Z. Q. (2018). Phenolic and flavonoid contents of mandarin (*Citrus reticulata* Blanco) fruit tissues and their antioxidant capacity as evaluated by DPPH and ABTS methods. *Journal of Integrative Agriculture*, *17*(1), 256-263.

Appendix

S/N	Sample	Saponin (%)	Tannin (%)	Alkaloid (%)	Flavonoid (%)	Oxalate (%)
1	<i>Xanthosoma Sagittifolium</i>	0.625±0.05	0.0000985±0.00	0.17±0.00	2.76±0.25	0.558±0.01
2	<i>Colocasia Esculenta</i>	0.682±0.00	0.0001235±0.05	1.065±0.00	1.64±0.00	0.3685±0.05

Table 1: Phytochemical profiles of *Xanthosoma Sagittifolium* and *Colocasia Esculenta*